



Section 24(c) Special Local Need Label

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

### RESTRICTED USE PESTICIDE Due to Acute Toxicity

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

## GRAMOXONE<sup>®</sup> MAX

EPA REG. NO. 100-1074

EPA SLN No. ID-010008

### After Planting – Preemergence Treatment for Potatoes

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Gramoxone Max Herbicide label.

This label must be in the possession of the user at the time of application.

**POTATOES (AFTER PLANTING-PREEMERGENCE TREATMENT).** Aerial application. Apply 11 to 20 fluid oz. per acre in 5 to 10 gallons of water. Add a nonionic surfactant containing at least 50% surface active ingredient at 1-2 pints per 100 gallons of dilute spray. The application should be delayed to provide maximum weed and grass emergence but should be applied not later than ground cracking, before potatoes have emerged. Application made after crop emergence has reduced yields of certain sensitive varieties. Weeds and grasses emerging after application will not be controlled. Postemergence cultivations should be performed as required to control subsequent weed and grass emergence.

**CHEMIGATION:** Do not apply this product through any type of irrigation system.

24(c) registrant:  
Syngenta Crop Protection, Inc.  
P.O. Box 18300  
Greensboro, NC 27419-8300

EPA SLN No. ID-010008

Label Code:

Word - IDGramoxoneMaxPotatoesSLNLbl